

2021-07-01

EFFECTS OF RISING CO₂ ON ALGAE IN JAPAN

Hall-Spencer, Jason

<http://hdl.handle.net/10026.1/17742>

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

This is the author's accepted manuscript. The final published version of this work (the version of record) was published by Taylor & Francis in *Phycologia* online on 5 August 2021 at <https://www.tandfonline.com/doi/pdf/10.1080/00318884.2021.1922050>

This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

Effects of rising CO₂ on algae in Japan

Jason Hall-Spencer¹, Sylvain Agostini², Ben Harvey², Lucia Porzio Shigeki Wada²

¹School of Biological and Marine Sciences, University of Plymouth, UK

² Shimoda Marine Research Center, University of Tsukuba, Japan

Shimoda Marine Research Centre has set up a platform for international research to assess the benefits of reducing multiple stressors for sustainable development of the Blue Economy. A closely monitored gradient in seawater CO₂ is available to assess ocean acidification, deoxygenation, eutrophication, warming and harmful species. We welcome collaborations as CO₂ seep show profound effects of acidification on micro- and macroalgae, as well as the ecological tipping points that drive shifts in coastal systems¹.

Recent sea surface warming has killed-off kelp forests around Shikine island. Here, coralline algae decline as CO₂ levels rise from 300 ppm to 900 ppm near seeps and perennial macroalgae are replaced by opportunistic turf algae and summer diatom blooms². Benthic carbon fixation is increased at high CO₂ - but most of it is exported offshore and the resilience of coastal communities to typhoons is diminished³. In the UN Decade of Ocean Science, data from this field facility will be used help inform people that need to adapt to global changes affecting aquaculture, fisheries and coastal tourism.

Financing: 'International Educational and Research Laboratory Program', University of Tsukuba. JSPS KAKENHI Grant 17K17622, Ministry of Environment, Japan (Suishinhi: 4RF-1701)

Keywords: Global change, marine ecology, CO₂ seeps

References

1. Hall-Spencer J.M., Harvey B.P. (2019) Ocean acidification impacts on coastal ecosystem services due to habitat degradation. *Emerging Topics in Life Sciences* 3 197–206
2. Harvey, B.P. et al. (*in press*) Feedback mechanisms stabilise degraded turf algal systems at a CO₂ seep site. *Communications Biology*.
3. Wada, S. et al. (*in press*). Ocean acidification increases phytobenthic carbon fixation and export in a warm-temperate system. *Estuarine, Coastal and Shelf Science*